



Minnesota Multiphasic  
Personality Inventory-2  
Restructured Form™

---

## Interpretive Report: Clinical Settings

---

MMPI-2-RF®

Minnesota Multiphasic Personality Inventory-2-Restructured Form™

*Yossef S. Ben-Porath, PhD, & Auke Tellegen, PhD*

---

ID Number: 1a  
Age: 26  
Gender: Male  
Marital Status: Married  
Years of Education: 14  
Date Assessed: 08/02/2008

**PEARSON**

**PsychCorp**


Copyright © 2008 by the Regents of the University of Minnesota. All rights reserved.  
Distributed exclusively under license from the University of Minnesota by NCS Pearson, Inc. Portions reproduced from the MMPI-2-RF test booklet. Copyright © 2008 by the Regents of the University of Minnesota. All rights reserved. Portions excerpted from the *MMPI-2-RF Manual for Administration, Scoring, and Interpretation*. Copyright © 2008 by the Regents of the University of Minnesota. All rights reserved.

**MMPI-2-RF** and the **MMPI-2-RF logo** are registered trademarks and **Minnesota Multiphasic Personality Inventory-2-Restructured Form** is a trademark of the University of Minnesota. **Pearson**, the **PSI logo**, and **PsychCorp** are trademarks in the U.S. and/or other countries of Pearson Education, Inc., or its affiliate(s).

### TRADE SECRET INFORMATION

Not for release under HIPAA or other data disclosure laws that exempt trade secrets from disclosure.

### MMPI-2-RF Validity Scales



Raw Score:	3	12	2	0	0	6	4	9
T Score:	48	57 T	51	42	42	45	57	55
Response %:	100	100	100	100	100	100	100	100
Cannot Say (Raw):	0							Percent True (of items answered): 40%


Comparison Group Data: MMPI-2-RF Normative (Men), N = 1,138

Mean Score (◇--◇):	50	50	50	51	50	48	50	50
Standard Dev (±1 SD):	11	11	10	10	10	9	10	10

The highest and lowest T scores possible on each scale are indicated by a "---"; MMPI-2-RF T scores are non-gendered.

VRIN-r	Variable Response Inconsistency	Fs	Infrequent Somatic Responses
TRIN-r	True Response Inconsistency	FBS-r	Symptom Validity
F-r	Infrequent Responses	L-r	Uncommon Virtues
Fp-r	Infrequent Psychopathology Responses	K-r	Adjustment Validity

### MMPI-2-RF Higher-Order (H-O) and Restructured Clinical (RC) Scales



Raw Score:	4	1	6	4	1	1	6	4	4	4	1	9
T Score:	41	48	50	51	42	38	49	49	70	46	47	45
Response %:	100	100	100	100	100	100	100	100	100	100	100	100


Comparison Group Data: MMPI-2-RF Normative (Men), N = 1,138

Mean Score (◊---◊):	49	50	53	49	49	50	51	52	51	48	50	51
Standard Dev (±1 SD):	10	10	10	9	9	10	10	10	10	9	10	11

The highest and lowest T scores possible on each scale are indicated by a "---"; MMPI-2-RF T scores are non-gendered.

- |  |                           |                                     |
|--|---------------------------|-------------------------------------|
| EID Emotional/Internalizing Dysfunction  | RCd Demoralization        | RC6 Ideas of Persecution            |
| THD Thought Dysfunction                  | RC1 Somatic Complaints    | RC7 Dysfunctional Negative Emotions |
| BXD Behavioral/Externalizing Dysfunction | RC2 Low Positive Emotions | RC8 Aberrant Experiences            |
|  | RC3 Cynicism              | RC9 Hypomanic Activation            |
|  | RC4 Antisocial Behavior   |                                     |

### MMPI-2-RF Somatic/Cognitive and Internalizing Scales



Raw Score:	1	0	0	0	2	0	1	0	2	1	0	1	0	3
T Score:	46	46	42	41	54	45	52	42	48	43	44	47	43	48
Response %:	100	100	100	100	100	100	100	100	100	100	100	100	100	100


Comparison Group Data: MMPI-2-RF Normative (Men), N = 1,138

Mean Score (◇--◇):	50	49	48	50	50	49	50	49	49	49	49	50	48	46
Standard Dev (±1 SD):	9	9	9	10	10	10	10	9	9	9	9	10	9	8

The highest and lowest T scores possible on each scale are indicated by a "---"; MMPI-2-RF T scores are non-gendered.

MLS	Malaise	SUI	Suicidal/Death Ideation	AXY	Anxiety
GIC	Gastrointestinal Complaints	HLP	Helplessness/Hopelessness	ANP	Anger Proneness
HPC	Head Pain Complaints	SFD	Self-Doubt	BRF	Behavior-Restricting Fears
NUC	Neurological Complaints	NFC	Inefficacy	MSF	Multiple Specific Fears
COG	Cognitive Complaints	STW	Stress/Worry		

### MMPI-2-RF Externalizing, Interpersonal, and Interest Scales



Raw Score:	2	0	0	2	1	1	1	3	0	0	7
T Score:	57	41	37	44	44	39	43	50	44	33	69
Response %:	100	100	100	100	100	100	100	100	100	100	100


Comparison Group Data: MMPI-2-RF Normative (Men), N = 1,138

Mean Score (◇--◇):	52	52	51	49	49	49	51	49	51	47	56
Standard Dev (±1 SD):	11	10	11	10	10	10	10	9	11	10	9

The highest and lowest T scores possible on each scale are indicated by a "---"; MMPI-2-RF T scores are non-gendered.

JCP	Juvenile Conduct Problems	FML	Family Problems	AES	Aesthetic-Literary Interests
SUB	Substance Abuse	IPP	Interpersonal Passivity	MEC	Mechanical-Physical Interests
AGG	Aggression	SAV	Social Avoidance		
ACT	Activation	SHY	Shyness		
		DSF	Disaffiliativeness		

### MMPI-2-RF PSY-5 Scales



Raw Score:	12	1	7	2	3
T Score:	60	47	51	40	42
Response %:	100	100	100	100	100

Comparison Group Data: MMPI-2-RF Normative (Men), N = 1,138

Mean Score (◇--◇):	52	50	54	49	51
Standard Dev (±1 SD):	11	10	10	10	11

The highest and lowest T scores possible on each scale are indicated by a "---"; MMPI-2-RF T scores are non-gendered.

- AGGR-r Aggressiveness-Revised
- PSYC-r Psychoticism-Revised
- DISC-r Disconstraint-Revised
- NEGE-r Negative Emotionality/Neuroticism-Revised
- INTR-r Introversion/Low Positive Emotionality-Revised

---

*This interpretive report is intended for use by a professional qualified to interpret the MMPI-2-RF. The information it contains should be considered in the context of the test taker's background, the circumstances of the assessment, and other available information.*

---

## **SYNOPSIS**

This is a valid MMPI-2-RF protocol. Scores on the substantive scales indicate thought dysfunction. Dysfunctional thinking findings relate to ideas of persecution.

## **PROTOCOL VALIDITY**

This is a valid MMPI-2-RF protocol. There are no problems with unscorable items. The test taker responded to the items relevantly on the basis of their content, and there are no indications of over- or under-reporting.

## **SUBSTANTIVE SCALE INTERPRETATION**

*Clinical symptoms, personality characteristics, and behavioral tendencies of the test taker are described in this section and organized according to an empirically guided framework. Statements containing the word "reports" are based on the item content of MMPI-2-RF scales, whereas statements that include the word "likely" are based on empirical correlates of scale scores. Specific sources for each statement can be viewed with the annotation features of this report.*

### **Somatic/Cognitive Dysfunction**

There are no indications of somatic or cognitive dysfunction in this protocol.

### **Emotional Dysfunction**

There are no indications of emotional-internalizing dysfunction in this protocol. The test taker reports a high level of psychological wellbeing, a wide range of emotionally positive experiences, and feeling confident and energetic.

### **Thought Dysfunction**

The test taker reports significant persecutory ideation such as believing that others seek to harm him. He is likely to be suspicious of and alienated from others, to experience interpersonal difficulties as a result of suspiciousness, and to lack insight.

### **Behavioral Dysfunction**

There are no indications of maladaptive externalizing behavior in this protocol. The test taker reports a below average level of aggressive behavior.

### **Interpersonal Functioning Scales**

These scales provide no further evidence of dysfunction.

### **Interest Scales**

The test taker reports an above average number of interests in activities or occupations of a mechanical or physical nature (e.g., fixing and building things, the outdoors, sports). Individuals who respond in this manner are likely to be adventure- and sensation-seeking. He indicates little or no interest in activities or occupations of an aesthetic or literary nature (e.g., writing, music, the theater).

## **DIAGNOSTIC CONSIDERATIONS**

*This section provides recommendations for psychodiagnostic assessment based on the test taker's MMPI-2-RF results. It is recommended that he be evaluated for the following:*

### **Thought Disorders**

- Disorders involving persecutory ideation

## **TREATMENT CONSIDERATIONS**

*This section provides inferential treatment-related recommendations based on the test taker's MMPI-2-RF scores.*

### **Psychotherapy Process Issues**

- Persecutory ideation may interfere with forming a therapeutic relationship and treatment compliance.

### **Possible Targets for Treatment**

- Persecutory ideation

## **ITEM-LEVEL INFORMATION**

### **Unscorable Responses**

The test taker produced scorable responses to all the MMPI-2-RF items.

## Critical Responses

Seven MMPI-2-RF scales--Suicidal/Death Ideation (SUI), Helplessness/Hopelessness (HLP), Anxiety (AXY), Ideas of Persecution (RC6), Aberrant Experiences (RC8), Substance Abuse (SUB), and Aggression (AGG)--have been designated by the test authors as having critical item content that may require immediate attention and follow-up. Items answered by the individual in the keyed direction (True or False) on a critical scale are listed below if his T score on that scale is 65 or higher. The percentage of the MMPI-2-RF normative sample (NS) and of the MMPI-2-RF Normative (Men) comparison group (CG) that answered each item in the keyed direction are provided in parentheses following the item content.

Ideas of Persecution (RC6, T Score = 70)

- 194. Item Content Omitted (True; NS 17.1%, CG 17.7%)
- 212. Item Content Omitted (False; NS 9.1%, CG 11.2%)
- 233. Item Content Omitted (True; NS 5.5%, CG 5.3%)
- 310. Item Content Omitted (True; NS 3.0%, CG 3.3%)



### Special Note:

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

## End of Report

This and previous pages of this report contain trade secrets and are not to be released in response to requests under HIPAA (or any other data disclosure law that exempts trade secret information from release). Further, release in response to litigation discovery demands should be made only in accordance with your profession's ethical guidelines and under an appropriate protective order.