



# MBMD° Pain Patient Reports

#### MILLON® BEHAVIORAL MEDICINE DIAGNOSTIC

Theodore Millon, PhD, DSc, Michael Antoni, PhD, Carrie Millon, PhD, Sarah Minor, PhD, and Seth Grossman, PsyD

## Two Pain Patient reports broaden clinical usefulness of the MBMD®

The MBMD Pain Patient Reports provide added breadth to a well-established instrument, helping clinicians and healthcare professionals identify psychosocial assets and liabilities that may affect an individual's response to treatment in a variety of pain settings.

This latest update to the MBMD is designed to help you more effectively manage care for pain patients, whether they are being evaluated for surgery or assessed for a treatment or rehabilitation program. In addition to the customizable Pain Patient Interpretive Report, a concise Profile Report is also available.

### Backed by extensive research to ensure relevance for pain settings

Developed by renowned test author Theodore Millon, PhD, DSc, the MBMD Pain Patient Reports are based on a comprehensive review of literature, input from experienced practitioners, and extensive reliability and validity data.

The normative sample comprises 1,200 patients in the United States who are being treated for issues involving chronic pain. Norm-referenced scores compare the patient to both a general medical norm sample and a chronic pain norm sample. A new manual supplement provides details on the development and use of the reports.

#### Relevant for multiple applications

The MBMD Pain Patient Reports are useful for a variety of purposes, including:

- Pre-treatment psychosocial evaluations to help professionals select suitable therapies
- General and expanded behavioral health evaluations to help determine the most appropriate intervention, such as individual psychotherapy, pain management groups, biofeedback, functional restoration/work hardening programs, or multidisciplinary pain programs
- Monitoring of treatment interventions and outcomes

ALWAYS LEARNING PEARSON

### **Customized reports for Pre-surgical** and Non-surgical patients

The Interpretive Report includes tailored considerations for clinicians working with pain patients to help serve as a guide in making prudent judgments. A three-page section in the report can be customized for either pre-surgical or nonsurgical populations.

- The Presurgical Pain Patient Report focuses on surgical risk factors and includes information on major and secondary outcome risks as well as postsurgical patient behavior.
- The Nonsurgical Pain Patient Report includes pre-treatment considerations, information on psychosocial management needs, and potential long-term gains and challenges.
- A one-page Healthcare Provider Summary section of the reports provides healthcare professionals with a useful and concise summary of the patient's potential assets and weaknesses.
- These population-specific reports are available with Q-global® web-based or Q Local™ desktop-based scoring and reporting software as well as with mail-in and fax-in scoring.

	010, Page 4	Sample Presurgical Repor
	PRESURGICAL PAIN PATIENT S	SUMMARY
reviews, cli	vizations in the following tables are credible and discriminating proba inical experience, and a few early empirical studies. As such, they sho o clinicians in making prudent and tentative judgments.	
I. PRESU	URGICAL CONSIDERATIONS	
The researchow they re	-Provider Communications th literature and clinical experience indicate that the interpersonal co- elate to others and the way they may relate to healthcare providers. Pro- when communicating with the patient as noted below.	roviders should consider the following
	The following healthcare orientation:	Is considered:
	Work to increase patient self-reliance Maintain strong focus on patient self-interest	Helpful ADVISABLE
	The following risk factor:	Is considered:
F	1. Depression	Moderate MARKED
		Moderate
	Depression     Anticipatory anxiety	Moderate MARKED
	Depression     Anticipatory anxiety     Cognitive deficits	Moderate MARKED low
	Depression     Anticipatory anxiety     Cognitive deficits     Pain sensitivity	Moderate MARKED low MARKED
	1. Depression 2. Anticipatory anxiety 3. Cognitive deficits 4. Pain sensitivity 5. Lack of social support	Moderate  MARKED  low  MARKED  low
	I. Depression  2. Anticipatory anxiety  3. Cognitive deficits  4. Pain sensitivity  5. Lack of social support  6. Medication abuse	Moderate  MARKED  low  MARKED  low  MARKED  low  Moderate
Although th	1. Depression 2. Anticipatory makey 3. Cognitive deficits 4. Pain sensitivity 5. Lack of social support 6. Medication also about 7. Problematic compliance 7. Problematic compliance 8. Calsaryphania 7. Surplice of Compliance 8. Calsaryphania 7. Surplice of Compliance 8. Calsaryphania 8. Calsaryphania 8. Surplice of Compliance 8. Surplice	Moderate MARKED Jor MARKED Jor MARKED Jor Moderate Jor Moderate Jor MARKED Jor Moderate Jor MARKED Jor Moderate Jor MARKED Jor MARKED Jor Moderate Jor MARKED Jor Moderate Jor MARKED Jor Moderate Jor MARKED Jor Moderate Jor Mod
Although th	1. Depression 2. Anticipatory survivy 3. Cognitive deficits 4. Palm sensitivity 5. Lack of social support 6. Medication abuses 7. Problematic compliance 7. Problematic compliance 8. Catterrophysics 9. Catterrophysics 9. Catterrophysics 9. Surgical Outcome Risks for characteristics limited below have not been studied to the extent that	Moderate MARKED low MARKED low MARKED low Moderate low MARKED they Moderate low MARKED they MARKED
Although th	1. Depression 2. Anticipatory smixty 3. Cognitive deficits 4. Palm ensimitive 5. Lack of social support 6. Medication abuse 7. Problematic compliance 7. Problematic compliance 8. Contamplating 197 Surgical Outstorm Risks 197 Surgical Outstorm Risks 198 Indian to the previous section, clinical experience suggests that they it listed in the previous section, clinical experience suggests that they it The following risk factors:	Marketa MARKED low MARKED low MAGENTE low Moderate low MARKED low Moderate low MARKED low Marketal low Marketal low Marketal low Marketal they can be included among the consensual sy indicate a risk for poor surgical outcome to considered:
Although th	1. Depression 2. Anticipatory makey 3. Cognitive deficits 4. Pen sensitivity 5. Lack of social support 6. Medication above 7. Problematic compliance 8. Calsaryphania 7. Surphile Conference of the Compliance 8. Calsaryphania 7. Surphile Conference State 8. Calsaryphania 8. Calsary	Moderate  MARKED  See  MARKED  Dee  MARKED  Dee  Moderate  low  MARKED  How  MARKED  Low  MARKED  MARKED  MARKED  MARKED  Low  MARKED  MARKED  MARKED  Low  MARKED  MARKED  MARKED  MARKED  MARKED  Low  MARKED  MARKED  MARKED  Low  MARKED  M
Although th	1. Depression 2. Anticipatory smixty 3. Cognitive deficits 4. Palm sensitivity 5. Lack of social support 6. Medication abuse 7. Problematic compliance 7. Problematic compliance 8. Contampatics 1. Problematic compliance 1. Increased in the previous section, clinical experience suggests that they are the compliance of the c	Moderate MARKED Low MARKED Low MARKED Low Moderate Low Moderate Low Moderate Low Moderate Low MARKED Low MARKED Low Low MARKED Low Low Low Low MARKED Low MARKED Low MARKED Low MARKED Low MARKED Low MARKED
Although th	1. Depression 2. Anticipatory makey 3. Cognitive deficits 4. Pain ensimity 5. Last of social support 6. Medication above 7. Pendematic compliance 8. Catastrophizing 7. Pendematic compliance 8. Catastrophizing 7. Surpliance 8. Catastrophizing 7. Line of the previous section, clinical experience suggests that they is 7. The following risk factor: 1. Line uself-carattrophization 2. Faz of illness complications 3. Self-indiagnees 8. Self	Moderate MARKED MSRED MSRED MSRED MSRED MSRED MSRED MSRED MSRED MSSRED MSSRED MARKED
Although th	1. Depression 2. Anticipatory smixty 3. Cognitive deficits 4. Palls neutritive 5. Lack of social support 6. Medication abuse 7. Production compliance 7. Production compliance 8. Contamorphism of the social support 1. Production of the social support 1. Interval to the previous section, editinal experience suggests that they are social support to the so	Moderate MARKED Low MARKED Low MARKED Low Moderate Low Moderate Low Moderate Low Moderate Low Moderate Low MARKED Low
Although th	1. Depression 2. Anticipatory makey 3. Cognitive deficits 4. Pain constitivity 5. Last of social support 6. Modication above 7. Problematic compliance 7. Problematic compliance 8. Cettasrophizing ary Surgical Outsome Risks he characteristics listed below have not been studied to the extent that listed in the previous section, clinical experience suggests that they is The following risk factor: 1. Low self-caracteristics and 5. Perior of illness complications 3. Self-indulgence 4. Oppositional attitude 5. Irrinability/hostingly	Moderate MARKED Mor MARKED Mor MARKED Mor Moderate Mor MARKED Mor MARKED MARKED MORE MORE MORE MORE MORE MORE MORE MORE
Although th	1. Depression 2. Anticipatory smixty 3. Cognitive deficits 4. Palm sensitivity 5. Lack of social support 6. Medication aboute 7. Problematic compliance 7. Problematic compliance 8. Contamorphisms 9. Contamorphi	Moderate MARKID Low MARKID Low MARKID Low Moderate Low MARKID Low Moderate Low MARKID Low Markin Mark

#### **About the Authors**

**Theodore Millon, PhD, DSc,** is a leading psychological theorist, renowned for his APA-award-winning work on an evolutionary theory of personality and psychopathology. He has been a full professor at Harvard Medical School, the University of Illinois, and the University of Miami. Principal author of the Millon™ Inventories, Dr. Millon has written or edited more than 30 books, and has contributed more than 250 chapters and articles to numerous books and journals in the field. In the past decade, he has received psychology's three highest commendations, the APA Presidential Citation, the APA's distinguished Professional Contribution to Research, and the APF's Gold Lifetime Achievement Award. With support from colleagues and Pearson, Dr. Millon established the Institute for Advanced Studies in Personology and Psychopathology, which he directs as Dean and Scientific Director.

Michael H. Antoni, PhD, is professor of Psychology and Psychiatry and Behavioral Sciences at the University of Miami, leads the Biobehavioral Oncology Program at the Sylvester Comprehensive Cancer Center, and is a licensed clinical psychologist in Florida. His research focuses on psychoneuroimmunology applied to viral infections and certain human cancers. He has led multiple NIH-funded efforts testing the effects of cognitive behavioral stress management (CBSM) interventions on psychological adjustment, immune system functioning and health outcomes in HIV-infected men and women, women with Human Papilloma Virus-associated cervical dysplasia, women with breast cancer, men with prostate cancer, and men and women with chronic fatigue syndrome. He is a fellow of the Society of Behavioral Medicine, and is associate editor for the International Journal of Behavioral Medicine and Psychology and Health. He has published more than 400 journal articles, abstracts, chapters, and books in the area of health psychology.

Visit PearsonClinical.com/MBMD or call 800.627.7271 for more information.

800.627.7271 | **Psych**Corp

PearsonClinical.com

Copyright © 2017 Pearson Education. All rights reserved. Pearson, Q-global, Q Local, PSI design, and PsychCorp are trademarks, in the U.S. and/or other countries, of Pearson Education, Inc. or its affiliates. Millon and MBMD are registered trademarks of DICANDRIEN, Inc. CLINA15777-3044 SR 6/17

**PEARSON** ALWAYS LEARNING