



Minnesota Multiphasic
Personality Inventory-2
Restructured Form®

MMPI-2-RF® Bibliography

Revised September 2012

MMPI-2-RF References by Topic

It is the publisher's intention to provide test users a current and comprehensive list of references to publications on the MMPI-2-RF. Therefore, this list will be updated as new publications appear. Please note that publications on the RC Scales, introduced in 2003, relate to the MMPI®-2 as well as to the MMPI-2-RF.

Basic Sources on the MMPI-2-RF

- Ben-Porath, Y. S. (2012). *Interpreting the MMPI-2-RF*. Minneapolis: University of Minnesota Press.
- Ben-Porath, Y. S., & Tellegen, A. (2008/2011). *MMPI-2-RF (Minnesota Multiphasic Personality Inventory-2 Restructured Form): Manual for administration, scoring, and interpretation*. Minneapolis: University of Minnesota Press.
- Ben-Porath, Y. S., & Tellegen, A. (2011). *MMPI-2-RF (Minnesota Multiphasic Personality Inventory-2 Restructured Form): User's guide for reports (2nd Edition)*. Minneapolis: University of Minnesota Press.
- Tellegen, A., & Ben-Porath, Y. S. (2008/2011). *MMPI-2-RF (Minnesota Multiphasic Personality Inventory-2 Restructured Form): Technical manual*. Minneapolis: University of Minnesota Press.
- Tellegen, A., Ben-Porath, Y. S., McNulty, J. L., Arbisi, P. A., Graham, J. R., & Kaemmer, B. (2003). *MMPI-2 Restructured Clinical (RC) Scales: Development, validation, and interpretation*. Minneapolis: University of Minnesota Press.

Publications on the MMPI-2-RF

General

- Archer, R. P. (2006). A perspective on the Restructured Clinical (RC) Scale project. *Journal of Personality Assessment*, 87, 179–185. doi: [10.1207/s15327752jpa8702_07](https://doi.org/10.1207/s15327752jpa8702_07)
- Ben-Porath, Y. S. (2006). Differentiating normal and abnormal personality with the MMPI-2. In S. Strack (Ed.), *Differentiating normal and abnormal personality* (2nd ed., pp. 337–381). NY: Springer.
- Ben-Porath, Y. S., & Tellegen, A. (2008). Empirical correlates of the MMPI-Restructured Clinical (RC) Scales in mental health, forensic, and non-clinical settings: An introduction. *Journal of Personality Assessment*, 90, 119–121. doi: [10.1080/00223890701845120](https://doi.org/10.1080/00223890701845120)
- Bolinski, P. K., & Nichols, D. S. (2011). Construct drift in the MMPI-2 Restructured Clinical Scales: Further evidence and a possible historic example. *Journal of Clinical Psychology*, 67, 1–11. doi: [10.1002/jclp.20814](https://doi.org/10.1002/jclp.20814)
- Burchett, D. L., & Ben-Porath, Y. S. (2010). The Impact of Overreporting on MMPI-2-RF Substantive Scale Score Validity. *Assessment*, 17, 497–516. doi: [10.1177/1073191110378972](https://doi.org/10.1177/1073191110378972)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Caldwell, A. B. (2006). Maximal measurement or meaningful measurement: The interpretive challenges of the MMPI-2 Restructured Clinical (RC) Scales. *Journal of Personality Assessment*, 87, 193–201. doi: [10.1207/s15327752jpa8702_09](https://doi.org/10.1207/s15327752jpa8702_09)
- Cox, A., Pant, H., Gilson, A. N., Rodriguez, J. S., Young, K. R., Kwon, S., & Weed, N. C., (2012). Effects of augmenting response options on MMPI-2 RC Scale psychometrics. *Journal of Personality Assessment*, 94, 613–619. doi: [10.1080/00223891.2012.700464](https://doi.org/10.1080/00223891.2012.700464)
- Eaton, N., Krueger, R. F., South, S. C., Simms, L. J. (2011). Contrasting prototypes and dimensions in the classification of personality pathology: Evidence that dimensions, but not prototypes, are robust. *Psychological Medicine*, 41, 1151–1163. doi: [10.1017/S0033291710001650](https://doi.org/10.1017/S0033291710001650)
- Finn, S. E., & Kamphuis, J. H. (2006). The MMPI-2 RC Scales and restraints to innovation, or “What have they done to my song?” *Journal of Personality Assessment*, 87, 202–210. doi: [10.1207/s15327752jpa8702_10](https://doi.org/10.1207/s15327752jpa8702_10)
- Forbey, J. D., & Ben-Porath, Y. S. (2007). Computerized adaptive personality testing: A review and illustration with the MMPI-2 Computerized Adaptive Version (MMPI-2-CA). *Psychological Assessment*, 19, 14–24. doi: [10.1037/1040-3590.19.1.14](https://doi.org/10.1037/1040-3590.19.1.14)
- Graham, J. R. (2011). *MMPI-2: Assessing Personality and Psychopathology (5th edition)*. New York: Oxford University Press.
- Greene, R. L. (2011). *MMPI-2/MMPI-2-RF: An Interpretive Manual (Third Edition)*. New York: Allyn & Bacon.
- Greene, R. L., Rouse, S. V., Butcher, J. N., Nichols, D. S., & Williams, C. L. (2009). The MMPI-2 Restructured Clinical (RC) Scales and redundancy: Reply to Tellegen, Ben-Porath, and Sellbom. *Journal of Personality Assessment*, 91, 222–226. doi: [10.1080/00223890902800825](https://doi.org/10.1080/00223890902800825)
- Harkness, A. R., Finn, J. A., McNulty, J. L., & Shields, S. M. (2012). The Personality Psychopathology Five (PSY-5): Recent constructive replication and assessment literature review. *Psychological Assessment*, 24, 432–443. doi: [10.1037/a0025830](https://doi.org/10.1037/a0025830)
- Hoelzle, J. B., & Meyer, G. J. (2008). The factor structure of the MMPI-2 Restructured Clinical (RC) Scales. *Journal of Personality Assessment*, 90, 443–455. doi: [10.1080/00223890802248711](https://doi.org/10.1080/00223890802248711)
- Lee, T. C., Forbey, J. D., & Ritchey, K. A. (2011). The impact of emotional priming on MMPI-2 scale scores. *Assessment*, 18, 102–105. doi: [10.1177/1073191110392496](https://doi.org/10.1177/1073191110392496)
- Nichols, D. S. (2006a). The trials of separating bath water from baby: A review and critique of the MMPI-2 Restructured Clinical Scales. *Journal of Personality Assessment*, 87, 121–138. doi: [10.1207/s15327752jpa8702_02](https://doi.org/10.1207/s15327752jpa8702_02)
- Nichols, D. S. (2006b). Commentary on Rogers, Sewell, Harrison, & Jordan (2006). *Journal of Personality Assessment*, 87, 172–174. doi: [10.1207/s15327752jpa8702_05](https://doi.org/10.1207/s15327752jpa8702_05)
- Rogers, R., & Sewell, K. W. (2006). MMPI-2 at the crossroads: Aging technology or retrofitting. *Journal of Personality Assessment*, 87, 175–178. doi: [10.1207/s15327752jpa8702_06](https://doi.org/10.1207/s15327752jpa8702_06)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Rogers, R., Sewell, K. W., Harrison, K. S., & Jordan, M. J. (2006). The MMPI-2 Restructured Clinical Scales: A paradigmatic shift in scale development. *Journal of Personality Assessment*, 87, 139–147. doi: [10.1207/s15327752jpa8702_03](https://doi.org/10.1207/s15327752jpa8702_03)
- Rouse, S. V., Greene, R. L., Butcher, J. N., Nichols, D. S., & Williams, C. L. (2008). What do the MMPI-2 Restructured Clinical Scales reliably measure? Answers from multiple research settings. *Journal of Personality Assessment*, 90, 435–442. doi: [10.1080/00223890802248695](https://doi.org/10.1080/00223890802248695)
- Sellbom, M., Ben-Porath, Y. S., & Bagby, R. M. (2008). On the hierarchical structure of mood and anxiety disorders: Confirmatory evidence and elaboration of a model of temperament markers. *Journal of Abnormal Psychology*, 117, 576–590. doi: [10.1037/a0012536](https://doi.org/10.1037/a0012536)
- Sellbom, M., Ben-Porath, Y. S., Graham, J. R., Arbisi, P. A., & Bagby, R. M. (2005). Susceptibility of the MMPI-2 Clinical, Restructured Clinical (RC), and Content Scales to overreporting and underreporting. *Assessment*, 12, 79–85. doi: [10.1177/1073191104273515](https://doi.org/10.1177/1073191104273515)
- Simms, L. J. (2006). Bridging the divide: Comments on the Restructured Clinical Scales of the MMPI-2. *Journal of Personality Assessment* 87, 211–216. doi: [10.1207/s15327752jpa8702_11](https://doi.org/10.1207/s15327752jpa8702_11)
- Tellegen, A., Ben-Porath, Y. S., & Sellbom, M. (2009). Construct validity of the MMPI-2-Restructured Clinical (RC) Scales: Reply to Rouse, Greene, Butcher, Nichols, & Williams. *Journal of Personality Assessment*, 91, 211–221. doi: [10.1080/00223890902794192](https://doi.org/10.1080/00223890902794192)
- Tellegen, A., Ben-Porath, Y. S., Sellbom, M., Arbisi, P. A., McNulty, J. L., & Graham, J. R. (2006). Further evidence on the validity of the MMPI-2 Restructured Clinical (RC) Scales: Addressing questions raised by Rogers et al. and Nichols. *Journal of Personality Assessment*, 87, 148–171. doi: [10.1207/s15327752jpa8702_04](https://doi.org/10.1207/s15327752jpa8702_04)
- Van der Heijden, P. T., Egger, J. I. M., & Derkson, J. J. L. (2008). Psychometric evaluation of the MMPI-2 Restructured Clinical Scales in two Dutch Samples. *Journal of Personality Assessment*, 90, 456–464. doi: [10.1080/00223890802248745](https://doi.org/10.1080/00223890802248745)
- Van der Heijden, P. T., Egger, J. I. M., & Derkson, J. J. L. (2010). Comparability of scores on the MMPI-2-RF scales generated with the MMPI-2 and MMPI-2-RF booklets. *Journal of Personality Assessment*, 92, 254–259. doi: [10.1080/00223891003670208](https://doi.org/10.1080/00223891003670208)
- Viken, R. J., & Rose, R. J. (2007). Genetic variation and covariation in the original and Restructured Clinical Scales of the MMPI. *Journal of Abnormal Psychology*, 116, 842–847. doi: [10.1037/0021-843X.116.4.842](https://doi.org/10.1037/0021-843X.116.4.842)
- Weed, N. C. (2006). Syndromal complexity, paradigm shifts, and the future of validation research: A comment on Nichols and Rogers, Sewell, Harrison, & Jordan. *Journal of Personality Assessment*, 87, 217–222. doi: [10.1207/s15327752jpa8702_12](https://doi.org/10.1207/s15327752jpa8702_12)
- Weiss, P. A., Bell, K. J., & Weiss, W. U. (2010). Use of the MMPI-2 RC Scales to detect criminal malingering. *Journal of Police and Criminal Psychology*, 25, 49–55. doi: [10.1007/s11896-009-9056-9](https://doi.org/10.1007/s11896-009-9056-9)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

Validity Scales

- Dionysus, K., Denney, R. L., & Halfaker, D. A. (2011). Detecting negative response bias with the Fake Bad Scale, Response Bias Scale, Henry-Heilbronner Index of the Minnesota Multiphasic Personality Inventory-2. *Archives of Clinical Neuropsychology*, 26, 81–88. doi: [10.1093/arclin/acq](https://doi.org/10.1093/arclin/acq)
- Dragon, W. R., Ben-Porath, Y. S., & Handel, R. H. (2012). Examining the impact of unscorable item responses on the validity and interpretability of MMPI-2/MMPI-2-RF Restructured Clinical (RC) Scale scores. *Assessment*, 19, 101–113. doi: [10.1177/1073191111415362](https://doi.org/10.1177/1073191111415362)
- Gass, C. S., & Odland, A. P. (2012). Minnesota Multiphasic Personality Inventory-2 Revised Form [sic] Symptom Validity Scale-Revised (MMPI-2-RF FBS-r; also known as Fake Bad Scale): Psychometric characteristics in a nonlitigation neuropsychological setting. *Journal of Clinical and Experimental Neuropsychology*, 34, 561–570. doi: [10.1080/13803395.2012.666228](https://doi.org/10.1080/13803395.2012.666228)
- Gervais, R. O., Ben-Porath, Y. S., Wygant, D. B., & Green, P. (2007). Development and validation of a Response Bias Scale (RBS) for the MMPI-2. *Assessment*, 14, 196–208. doi: [10.1177/1073191106295861](https://doi.org/10.1177/1073191106295861)
- Gervais, R. O., Ben-Porath, Y. S., Wygant, D. B., & Green, P. (2008). Differential sensitivity of the Response Bias Scale and MMPI-2 Validity Scales to memory complaints. *The Clinical Neuropsychologist*, 22, 1061–1079. doi: [10.1080/13854040701756930](https://doi.org/10.1080/13854040701756930)
- Gervais, R. O., Ben-Porath, Y. S., & Wygant, D. B. (2009). Empirical correlates and interpretation of the MMPI-2-RF Cognitive Complaints (COG) scale. *The Clinical Neuropsychologist*, 23, 996–1015. doi: [10.1080/13854040902748249](https://doi.org/10.1080/13854040902748249)
- Gervais, R. O., Ben-Porath, Y. S., Wygant, D. B., & Sellbom, M. (2010). Incremental validity of the MMPI-2-RF over-reporting scales and RBS in assessing the veracity of memory complaints. *Archives of Clinical Neuropsychology*, 25, 274–284. doi: [10.1093/arclin/acq018](https://doi.org/10.1093/arclin/acq018)
- Gervais, R. O., Wygant, D. B., Sellbom, M., & Ben-Porath, Y. S. (2011). Associations between Symptom Validity Test failure and scores on the MMPI-2-RF validity and substantive scales. *Journal of Personality Assessment*, 93, 508–517. doi: [10.1080/00223891.2011.594132](https://doi.org/10.1080/00223891.2011.594132)
- Handel, R. W., Ben-Porath, Y. S., Tellegen, A., & Archer, R. P. (2010). Psychometric functioning of the MMPI-2-RF VRIN-r and TRIN-r Scales with varying degrees of randomness, acquiescence, and counter-acquiescence. *Psychological Assessment*, 22, 87–95. doi: [10.1037/a0017061](https://doi.org/10.1037/a0017061)
- Harp, J. P., Jasinski, L. J., Shandrea-Ochsner, A., Mason, L. H., & Berry, D. T. R. (2011). Detection of malingered ADHD using the MMPI-2-RF. *Psychological Injury and Law*, 4, 32–43. doi: [10.1007/s12207-011-9100-9](https://doi.org/10.1007/s12207-011-9100-9)
- Jones, A., & Ingram, M. V. (2011). A comparison of selected MMPI-2 and MMPI-2-RF Validity Scales in assessing effort on cognitive tests in a military sample. *The Clinical Neuropsychologist*, 25, 1207–1227. doi: [10.1080/13854046.2011.600726](https://doi.org/10.1080/13854046.2011.600726)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Jones, A., Ingram, M. V., & Ben-Porath, Y. S. (2012). Scores on the MMPI-2-RF scales as a function of increasing levels of failure on cognitive symptom validity tests in a military sample. *The Clinical Neuropsychologist*, 26, 790–815. doi: [org/10.1080/13854046.2012.693202](https://doi.org/10.1080/13854046.2012.693202)
- Lange, R. T., Sullivan, K. A., & Scott, C. (2010). Comparison of MMPI-2 and PAI validity indicators to detect feigned depression and PTSD symptom reporting. *Psychiatry Research*, 176, 229–235. doi: [10.1016/j.psychres.2009.03.004](https://doi.org/10.1016/j.psychres.2009.03.004)
- Marion, B. E., Sellbom, M., & Bagby, R. M. (2011). The detection of feigned psychiatric disorders using the MMPI-2-RF overreporting Validity Scales: An analog Investigation. *Psychological Injury and Law*, 4, 1–12. doi: [10.1007/s12207-011-9097-0](https://doi.org/10.1007/s12207-011-9097-0)
- Marion, B. E., Sellbom, M., Salekin, R. T., Toomey, J. A., Kucharski, T., & Duncan, S. (in press). An examination of the association between psychopathy and dissimulation using the MMPI-2-RF Validity Scales. *Law and Human Behavior*.
- Nelson, N. W., Sweet, J. J., & Heilbronner, R. L. (2007). Examination of the new MMPI-2 Response Bias Scale (Gervais): relationship with MMPI-2 validity scales. *Journal of Clinical and Experimental Neuropsychology*, 29, 67–72. doi: [10.1080/13803390500488546](https://doi.org/10.1080/13803390500488546)
- Rogers, R., Gillard, N., Berry, D. T. R., & Granacher, R. P. (2011). Effectiveness of the MMPI-2-RF Validity Scales for feigned mental disorders and cognitive impairment: A known-groups study. *Journal of Psychopathology and Behavioral Assessment*, 33, 355–367. doi: [10.1007/s10862-011-9222-0](https://doi.org/10.1007/s10862-011-9222-0)
- Schroeder, R. W., Baade, L. E., Peck, C. P., VonDran, E. ., Brockman, C. J., Webster, B. K., & Heinrichs, R. J. (2012). Validation of the MMPI-2-RF Validity Scales in criterion group neuropsychological samples. *The Clinical Neuropsychologist*, 26, 129–146. doi: [10.1080/13854046.2011.639314](https://doi.org/10.1080/13854046.2011.639314)
- Sellbom, M., & Bagby, R. M. (2008). Validity of the MMPI-2-RF (Restructured Form) L-r and K-r scales in detecting underreporting in clinical and nonclinical samples. *Psychological Assessment*, 20, 370–376. doi: [10.1037/a0012952](https://doi.org/10.1037/a0012952)
- Sellbom, M., & Bagby, R. M. (2010). The detection of over-reported psychopathology with the MMPI-2-RF (Restructured Form) Validity Scales. *Psychological Assessment*, 22, 757–767. doi: [10.1037/a0020825](https://doi.org/10.1037/a0020825)
- Sellbom, M., Toomey, A., Wygant, D., Kucharski, L. T., & Duncan, S. (2010). Utility of the MMPI-2-RF (Restructured Form) Validity Scales in detecting malingering in a criminal forensic setting: A known-groups design. *Psychological Assessment*, 22, 22–31. doi: [10.1037/a0018222](https://doi.org/10.1037/a0018222)
- Sellbom, M., Wygant, D. B., & Bagby, R. M. (2012). Utility of the MMPI-2-RF in detecting non-credible somatic complaints. *Psychiatry Research*, 197, 295–301. doi: [10.1016/j.psychres.2011.12.043](https://doi.org/10.1016/j.psychres.2011.12.043)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Smart, C. M., Nelson, N. W., Sweet, J. J., Bryant, F. B., Berry, D. T., Granacher, R. P., & Heilbronner, R. L. (2008). Use of MMPI-2 to predict cognitive effort: A hierarchically optimal classification tree analysis. *Journal of the International Neuropsychological Society*, 14, 842–852. doi: [10.1017/S1355617708081034](https://doi.org/10.1017/S1355617708081034)
- Sullivan, K. A., & Elliott, C. (2012). An investigation of the validity of the MMPI-2 Response Bias Scale using an analog simulation design. *The Clinical Neuropsychologist*, 26, 160–176. doi: [10.1080/13854046.2011.647084](https://doi.org/10.1080/13854046.2011.647084)
- Tolin, D. F., Steenkamp, M. M., Marx, B. P., & Litz, B. T. (2010). Detecting symptom exaggeration in combat veterans using the MMPI-2 symptom validity scales: A mixed group validation. *Psychological Assessment*, 22, 729–736. doi: [10.1037/a0020973](https://doi.org/10.1037/a0020973)
- Tsushima, W. T., Geling, O., & Fabriga, L. (2011). Comparison of MMPI-2 Validity Scale scores of personal injury litigants and disability claimants. *The Clinical Neuropsychologist*, 25, 1403–1414. doi: [10.1080/13854046.2011.613854](https://doi.org/10.1080/13854046.2011.613854)
- Van Dyke, S. A., Axelrod, B. N., & Schutte, C. (2010). The utility of the Post-conclusive Syndrome Questionnaire. *Archives of Clinical Neuropsychology*, 25, 634–639. doi: [10.1093/arclin/acq063](https://doi.org/10.1093/arclin/acq063)
- Whitney, K. A., Davis, J. J., Shephard, P. H., & Herman, S. M. (2008). Utility of the Response Bias Scale and other MMPI-2 validity scales in predicting TOMM performance. *Archives of Clinical Neuropsychology*, 23, 777–786. doi: [10.1016/j.acn.2008.09.001](https://doi.org/10.1016/j.acn.2008.09.001)
- Wygant, D. B., Anderson, J. L., Sellbom, M., Rapier, J. L., Algeier, L. M., & Granacher, R. P. (2011). Association of MMPI-2 Restructured Form (MMPI-2-RF) validity scales with structured malingering criteria. *Psychological Injury and Law*, 4, 13–23. doi: [10.1007/s12207-011-9098-z](https://doi.org/10.1007/s12207-011-9098-z)
- Wygant, D. B., Ben-Porath, Y. S., Arbisi, P. A., Berry, D. T. R., Freeman, D. B., & Heilbronner, R. L. (2009). Examination of the MMPI-2 Restructured Form (MMPI-2-RF) Validity Scales in civil forensic settings: Findings from simulation and known group samples. *Archives of Clinical Neuropsychology*, 27, 671–680. doi: [10.1093/arclin/acp073](https://doi.org/10.1093/arclin/acp073)
- Wygant, D. B., Sellbom, M., Gervais, R. O., Ben-Porath, Y. S., Stafford, K. P., Freeman, D. B., & Heilbronner, R. L. (2010). Further validation of the MMPI-2 and MMPI-2-RF Response Bias Scale: Findings from disability and criminal forensic settings. *Psychological Assessment*, 22, 745–756. doi: [10.1037/a0020042](https://doi.org/10.1037/a0020042)
- Young, J. C., & Gross, A. M. (2011). Detection of response bias and noncredible performance in adult Attention-Deficit/Hyperactivity Disorder. *Archives of Clinical Neuropsychology*, 26, 165–175. doi: [10.1093/arclin/acr013](https://doi.org/10.1093/arclin/acr013)
- Young, J. C., Kearns, L. A., & Roper, B. L. (2011). Validation of the MMPI-2 Response Bias Scale and Henry-Heilbronner Index in a U.S. veteran population. *Archives of Clinical Neuropsychology*, 3, 194–204. doi: [10.1093/arclin/acr015](https://doi.org/10.1093/arclin/acr015)
- Youngjohn, J. R., Weshba, R., Stevenson, M., Sturgeon, J., & Thomas, M. L. (2011). Independent validation of the MMPI-2-RF Somatic/Cognitive and Validity Scales in TBI litigants tested for effort. *The Clinical Neuropsychologist*, 25, 463–476. doi: [10.1080/13854046.2011.554444](https://doi.org/10.1080/13854046.2011.554444)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

Correctional Settings

- Clegg, C., Fremouw, W., Horacek, T., Cole, A., & Schwartz, R. (2010). Factors associated with treatment acceptance and compliance among incarcerated male sex offenders. *International Journal of Offender Therapy and Comparative Criminology*, 6, 880–897. doi: [10.1177/0306624X10376160](https://doi.org/10.1177/0306624X10376160).
- Forbey, J. D., Ben-Porath, Y. S., & Gartland, D. (2009). Validation of the MMPI-2 Computerized Adaptive Version (MMPI-2-CA) in a correctional intake facility. *Psychological Services*, 6, 279–292. doi: [10.1037/a0016195](https://doi.org/10.1037/a0016195)
- Kastner, R. M., Sellbom, M., & Lilienfeld, S. O. (2012). A comparison of the psychometric properties of the Psychopathic Personality Inventory Full-Length and Short-Form Versions. *Psychological Assessment*, 24, 261–267. doi: [10.1037/a0025832](https://doi.org/10.1037/a0025832)
- Megargee, Edwin I. (2006). *Using the MMPI-2 in Criminal Justice and Correctional Settings*. Minneapolis: University of Minnesota Press.
- Sellbom, M., Ben-Porath, Y. S., Patrick, C. J., Wygant, D. B., Gartland, D. M., & Stafford, K. P. (2012). Development and construct validation of MMPI-2-RF measures assessing global psychopathy, fearless-dominance, and impulsive-antisociality. *Personality Disorders: Theory, Research, and Treatment*, 3, 17–38. doi: [10.1037/a0023888](https://doi.org/10.1037/a0023888)

Forensic Settings

- Archer, E. M., Hagan, L. D., Mason, J., Handel, R. W., & Archer, R. P. (2012). MMPI-2-RF characteristics of custody evaluation litigants. *Assessment*, 19, 14–20. doi: [10.1177/1073191110397469](https://doi.org/10.1177/1073191110397469)
- Ben-Porath, Y. S. (in press). Addressing challenges to MMPI-2-RF-based testimony: Questions and answers. *Archives of Clinical Neuropsychology*.
- Downing, S. K., Denney, R. L., Spray, B. J., Houston, C. M., & Halfaker, D. H. (2008). Examining the relationship between the Restructured Scales and the Fake Bad Scale of the MMPI-2. *The Clinical Neuropsychologist*, 22, 680–688. doi: [10.1080/13854040701562825](https://doi.org/10.1080/13854040701562825)
- Gervais, R. O., Ben-Porath, Y. S., & Wygant, D. B. (2009). Empirical correlates and interpretation of the MMPI-2-RF Cognitive Complaints Scale. *The Clinical Neuropsychologist*, 23, 996–1015. doi: [10.1080/13854040902748249](https://doi.org/10.1080/13854040902748249)
- Henry, G. K., Heilbronner, R. L., Mittenberg, W., Enders, C., & Dombaski, K. (2009). Comparison of the MMPI-2 Restructured Demoralization Scale, Depression Scale, and Malingered Mood Disorder Scale in identifying non-credible symptom reporting in personal injury litigants and disability claimants. *The Clinical Neuropsychologist*, 23, 153–166. doi: [10.1080/13854040801969524](https://doi.org/10.1080/13854040801969524)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Henry, G. K., Heilbronner, R. L., Mittenberg, W., Enders, C., & Stanczal, S. R. (2008). Comparison of the Lees-Haley Fake Bad Scale, Henry-Heilbronner Index, and Restructured Clinical Scale I in identifying noncredible symptom reporting. *The Clinical Neuropsychologist*, 22, 919–929. doi: [10.1080/13854040701625853](https://doi.org/10.1080/13854040701625853)
- Mattson, C. A., Powers, B. K., Halfaker, D., Akenson, S. T., & Ben-Porath, Y. S. (2012). Predicting drug court completion with the MMPI-2-RF. *Psychological Assessment*, 24, 937–943. doi: [10.1037/a0028267](https://doi.org/10.1037/a0028267)
- Pinsoeneault, T. B., & Ezzo, F. R. (2012). A comparison of MMPI-2-RF profiles between child maltreatment and non-maltreatment custody cases. *Journal of Forensic Psychology Practice*, 12, 227–237. doi: [org/10.1080/15228932.2012.674469](https://doi.org/10.1080/15228932.2012.674469)
- Resendes, J., & Lecci, L. (2012). Comparing the MMPI-2 scale scores of parents involved in parental competency and child custody assessments. *Psychological Assessment*, 24, 1054–1059. doi: [10.1037/a0028585](https://doi.org/10.1037/a0028585)
- Rock, R. C., Sellbom, M., Ben-Porath, Y. S., & Salekin, R. T. (2012). Concurrent and predictive validity of psychopathy in a batterers intervention sample. *Law and Human Behavior*. doi: [10.1037/lhb0000006](https://doi.org/10.1037/lhb0000006)
- Sellbom, M. (2012). The MMPI-2-RF is ready for the Daubert challenge: Evidence, implications, and recommendations for use in court testimony. *Journal of Psychological Practice*, 17, 151–179.
- Sellbom, M. (2011). Elaborating on the construct validity of the Levenson Self-Report Psychopathy Scale in incarcerated and non-incarcerated samples. *Law and Human Behavior*, 35, 440–451. doi: [10.1007/s10979-010-9249-x](https://doi.org/10.1007/s10979-010-9249-x)
- Sellbom, M., & Bagby, R. M. (2009). Identifying PTSD personality subtypes in a workplace trauma sample. *Journal of Traumatic Stress*, 22, 471–475. doi: [10.1002/jts.20452](https://doi.org/10.1002/jts.20452)
- Sellbom, M., & Ben-Porath, Y. S. (2006). Forensic applications of the MMPI. In R. P. Archer (Ed.), *Forensic Uses of Clinical Assessment Instruments* (pp. 19–55). NJ: Lawrence Erlbaum Associates.
- Sellbom, M., Ben-Porath, Y. S., Baum, L. J., Erez, E., & Gregory, C. (2008). Predictive validity of the MMPI-2 Restructured Clinical (RC) Scales in a batterers' intervention program. *Journal of Personality Assessment*, 90, 129–135. doi: [10.1080/00223890701845153](https://doi.org/10.1080/00223890701845153)
- Sellbom, M., Ben-Porath, Y. S., & Stafford, K. P. (2007). A comparison of measures of psychopathic deviance in a forensic setting. *Psychological Assessment*, 19, 430–436. doi: [10.1037/1040-3590.19.4.430](https://doi.org/10.1037/1040-3590.19.4.430)
- Sellbom, M., Lee, T. T. C., Ben-Porath, Y. S., Arbisi, P. A., & Gervais, R. O. (2012). Differentiating PTSD symptomatology with the MMPI-2-RF (Restructured Form) in a forensic disability sample. *Psychiatry Research*, 197, 172–179. doi: [10.1016/j.psychres.2012.02.003](https://doi.org/10.1016/j.psychres.2012.02.003)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Stredny, R. V., Archer, R. P., & Mason, J. A. (2006). MMPI-2 and MCMI-III characteristics of parental competency examinees. *Journal of Personality Assessment*, 87, 113–115. doi: [10.1207/s15327752jpa8701_10](https://doi.org/10.1207/s15327752jpa8701_10)
- Thomas, M. L. & Youngjohn, J. R. (2009). Let's not get hysterical: Comparing the MMPI-2 Validity, Clinical, and RC Scales in TBI litigants tested for effort. *The Clinical Neuropsychologist*, 23, 1067–1084. doi: [10.1080/13854040902795000](https://doi.org/10.1080/13854040902795000)
- Vines, L. M., Wygant, D. B., & Gervais, R. O. (2012). Empirically guided case conceptualization of Posttraumatic Stress Disorder with the Minnesota Multiphasic Personality Inventory-2 Restructured Form (MMPI-2-RF) in a forensic disability evaluation. *Journal of Psychological Practice*, 17, 180–205.
- Wise, E. A. (2009). Selected MMPI-2 scores of forensic offenders in a community setting. *Journal of Forensic Psychology Practice*, 9, 299–309. doi: [10.1080/15228930902936048](https://doi.org/10.1080/15228930902936048)
- Wygant, D. B., & Sellbom, M. (in press). Viewing psychopathy from the perspective of the Personality Psychopathology Five model: Implications for DSM–5. *Journal of Personality Disorders*.
- Wygant, D. B., Sellbom, M., Ben-Porath, Y. S., Stafford, K. P., Freeman, D. B., & Heilbronner, R. I. (2007). The relation between symptom validity testing and MMPI-2 scores as a function of forensic evaluation context. *Archives of Clinical Neuropsychology*, 22, 488–499. doi: [10.1016/j.acn.2007.01.027](https://doi.org/10.1016/j.acn.2007.01.027)

Medical Settings

- Block, A. R., Ben-Porath, Y. S., & Marek, R. J. (in press). Psychological risk factors for poor outcome of spine surgery and spinal cord stimulator implant: A review of the literature and their assessment with the MMPI-2-RF. *The Clinical Neuropsychologist*.
- Block, A., Ohnmeiss, D., Ben-Porath, Y., & Burchett, D. (2011). Presurgical psychological screening: A new algorithm, including the MMPI-2-RF, for predicting surgery results. *Proceedings of the NASS 26th Annual Meeting/The Spine Journal 11 (2011) 1S–173S*. doi: [10.1016/j.spinee.2011.08.333](https://doi.org/10.1016/j.spinee.2011.08.333)
- Butcher, J. N., Hamilton, C. K., Rouse, S. V., & Cumella, E. J. (2006). The deconstruction of the Hy Scale of MMPI-2: Failure of RC3 in measuring somatic symptom expression. *Journal of Personality Assessment*, 87, 186–192. doi: [10.1207/s15327752jpa8702_08](https://doi.org/10.1207/s15327752jpa8702_08)
- D'Orzio, L. M., Meyerowitz, B. E., Korst, L. M., Romero, R., & Goodwin, T. M. (2011). Evidence against a link between Hyperemesis Gravidarum and personality characteristics from an ethnically diverse sample of pregnant women: A pilot study. *Journal of Women's Health*, 20, 137–143. doi: [10.1089/jwh.2009.1851](https://doi.org/10.1089/jwh.2009.1851)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Forbey, J. D., Arbisi, P. A., & Ben-Porath, Y. S. (2012). The MMPI-2 Computer Adaptive version (MMPI-2-CA) in a VA medical outpatient facility. *Psychological Assessment*, 24, 626–639. doi: [10.1037/a0026509](https://doi.org/10.1037/a0026509)
- Heinberg, L. J., Ashton, K., Windover, A. K., & Merrell, J. (2011). Older bariatric surgery candidates: Is there greater psychological risk than for young or midlife candidates? *Surgery for Obesity and Related Diseases*, 8, 616–622. doi: [10.1016/j.soard.2011.11.005](https://doi.org/10.1016/j.soard.2011.11.005)
- Locke, D. E. C., Kirlin, K. A., Thomas, M. L., Osborne, D., Hurst, D. F., Drazkowsi, J. F., Sirven, J. I., & Noe, K. H. (2010). The Minnesota Multiphasic Personality Inventory Restructured Form in the epilepsy monitoring unit. *Epilepsy and Behavior*, 17, 252–258. doi: [10.1016/j.yebeh.2009.12.004](https://doi.org/10.1016/j.yebeh.2009.12.004)
- Locke, D. E. C., Kirlin, K. A., Wershaba, R., Osborne, D., Drazkowski, J. F., Sirven, J. I., & Noe, K. H. (2011). Randomized comparison of the Personality Assessment Inventory and the Minnesota Multiphasic Personality Inventory-2, in the epilepsy monitoring unit. *Epilepsy and Behavior*, 21, 397–401. doi: [10.1016/j.yebeh.2011.05.023](https://doi.org/10.1016/j.yebeh.2011.05.023)
- Locke D. E. C., & Thomas M. L. (2011). Initial development of Minnesota Multiphasic Personality Inventory-2-Restructured Form (MMPI-2-RF) scales to identify patients with psychogenic non-epileptic seizures. *Journal of Clinical and Experimental Neuropsychology*, 33, 335–343. doi: [10.1080/13803395.2010.518141](https://doi.org/10.1080/13803395.2010.518141)
- McCord, D. M. & Drerup, L. C. (2011). Relative practical utility of the Minnesota Multiphasic Personality Inventory-2 Restructured Clinical Scales versus the Clinical Scales in a chronic pain patient sample. *Journal of Clinical and Experimental Neuropsychology*, 33, 140–146. doi: [10.1080/13803395.2010.495056](https://doi.org/10.1080/13803395.2010.495056)
- Merrell, J., Brethauer, S., Windover, A. K., Ashton, K., & Heinberg, L. J. (2012). Psychosocial correlates of pelvic floor disorders in women seeking bariatric surgery. *Surgery for Obesity and Related Diseases*, 8, 792–796. doi: [10.1016/j.soard.2012.01.019](https://doi.org/10.1016/j.soard.2012.01.019)
- Nelson, N. W., Hoelzle, J. B., McHuire, K. A., Sim, A. H., Goldman, D. J., Ferrier-Aurbach, A. G., Charlesworth, M. J., Arbisi, P. A., & Sponheim, S. R. (2011). Self-report of psychological function among OEF/OIF personnel who also report combat-related concussion. *The Clinical Neuropsychologist*, 25, 716–740. doi: [10.1080/13854046.2011.579174](https://doi.org/10.1080/13854046.2011.579174)
- Stroescu, I., Salinas, C. M., Nahab, F. B., & Stringer, A. Y. (2011). Long-term neurocognitive and neuroimaging outcomes in posterior reversible Encephalopathy Syndrome: Two case reports and implications. *The Clinical Neuropsychologist*, 25, 1386–1402. doi: [10.1080/13854046.2011.628339](https://doi.org/10.1080/13854046.2011.628339)
- Thomas, M. L., & Locke, D. E. C. (2010). Psychometric properties of the MMPI-2-RF Somatic Complaints (RC1) Scale. *Psychological Assessment*, 22, 492–503. doi: [10.1037/a0019229](https://doi.org/10.1037/a0019229)
- Wygant, D. B., Boutacoff, L. A., Arbisi, P. A., Ben-Porath, Y. S., Kelly, P. H., & Rupp, W. M. (2007). Examination of the MMPI-2 Restructured Clinical (RC) Scales in a sample of bariatric surgery candidates. *Journal of Clinical Psychology in Medical Settings*, 14, 197–205. doi: [10.1007/s10880-007-9073-8](https://doi.org/10.1007/s10880-007-9073-8)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

Mental Health Settings

- Arbisi, P. A., Erbes, C. R., Polusny, M. A., & Nelson, N. W. (2010). The concurrent and incremental validity of the Trauma Symptom Inventory in women reporting histories of sexual maltreatment. *Assessment*, *17*, 406–418. doi: [10.1177/107319110376163](https://doi.org/10.1177/107319110376163)
- Arbisi, P. A., Polusny, M. A., Erbes, C. R., Thusras, P., & Reddy, M. K. (2011). The Minnesota Multiphasic Personality Inventory-2 Restructured Form in National Guard soldiers screening positive for Posttraumatic Stress Disorder and Mild Traumatic Brain Injury. *Psychological Assessment*, *23*, 203–214. doi: [10.1037/a0021339](https://doi.org/10.1037/a0021339)
- Arbisi, P. A., Sellbom, M., & Ben-Porath, Y. S. (2008). Empirical correlates of the MMPI-2 Restructured Clinical (RC) Scales in psychiatric inpatients. *Journal of Personality Assessment*, *90*, 122–128. doi: [10.1080/00223890701845146](https://doi.org/10.1080/00223890701845146)
- Binford, A., & Liljequist, L. (2008). Behavioral correlates of selected MMPI-2 Clinical, Content, and Restructured Clinical scales. *Journal of Personality Assessment*, *60*, 608–614. doi: [10.1080/00223890802388657](https://doi.org/10.1080/00223890802388657)
- Bolinsky, P. K., Trumbetta, S. L., Hanson, D. R., & Gottesman, I. I. (2010). Predicting adult psychopathology from adolescent MMPIs: Some victories. *Personality and Individual Differences*, *49*, 324–330. doi: [10.1016/j.paid.2010.01.026](https://doi.org/10.1016/j.paid.2010.01.026)
- Castro, Y., Gordon, K. H., Brown, J. S., Cox, J. C., & Joiner, T. E. (2008). Examination of racial differences on the MMPI-2 Clinical and Restructured Clinical Scales in an outpatient sample. *Assessment*, *15*, 277–286. doi: [10.1177/1073191107312735](https://doi.org/10.1177/1073191107312735)
- Erbes, C. R., Polusny, M. A., Arbisi, P. A., & Koffel, E. (2012). PTSD symptoms in a cohort of National Guard Soldiers deployed to Iraq: Evidence for nonspecific and specific components. *Journal of Affective Disorders*. doi: [10.1016/j.jad.2012.05.013](https://doi.org/10.1016/j.jad.2012.05.013)
- Forbes, D., Elhai, J. D., Miller, M. W., & Creamer, M. (2010). Internalizing and externalizing classes in posttraumatic stress disorder: A latent class analysis. *Journal of Traumatic Stress*, *23*, 340–349. doi: [10.1002/jts.20526](https://doi.org/10.1002/jts.20526)
- Handel, R. W., & Archer, R. P. (2008). An investigation of the psychometric properties of the MMPI-2 Restructured Clinical (RC) Scales with mental health inpatients. *Journal of Personality Assessment*, *90*, 239–249. doi: [10.1080/00223890701884954](https://doi.org/10.1080/00223890701884954)
- Kamphuis, J. H., Arbisi, P. A., Ben-Porath, Y. S., & McNulty, J. L. (2008). Detecting comorbid Axis-II status among inpatients using the MMPI-2 Restructured Clinical Scales. *European Journal of Psychological Assessment*, *24*, 157–164. doi: [10.1027/1015-5759.24.3.157](https://doi.org/10.1027/1015-5759.24.3.157)
- McDevitt-Murphy, M. E., Weathers, F. W., Flood, A. M., Eakin, D. E., & Benson, T. A. (2007). The utility of the PAI and the MMPI-2 for discriminating PTSD, depression, and social phobia in trauma-exposed college students. *Assessment*, *14*, 181–195. doi: [10.1177/1073191106295914](https://doi.org/10.1177/1073191106295914)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Miller, M. W., Wolf, E. J., Harrington, K. M., Brown, T. A., Kaloupek, D. G., & Keane, T. M. (2010). An evaluation of competing models for the structure of PTSD symptoms using external measures of comorbidity. *Journal of Traumatic Stress*, 23, 631–638. doi: [10.1002/jts.20559](https://doi.org/10.1002/jts.20559)
- Ozonoff, S., Garcia, N., Clark, E., & Lainhart, J. E. (2005). MMPI-2 personality profiles of high-functioning adults with Autism Spectrum Disorders. *Assessment*, 12, 86–95. doi: [10.1177/1073191104273132](https://doi.org/10.1177/1073191104273132)
- Purdon, S. E., Purser, S. M., & Goddard, K. M. (2011). MMPI-2 Restructured Form over-reporting scales in first-episode psychosis. *The Clinical Neuropsychologist*, 25, 829–842. doi: [10.1080/13854046.2011.585141](https://doi.org/10.1080/13854046.2011.585141)
- Quilty, L. C., Sellbom, M., Tackett, J. L., & Bagby, R. M. (2009). Personality trait predictors of bipolar disorder: A replication and extension in a psychiatric sample. *Psychiatry Research*, 169, 159–163. doi: [10.1016/j.psychres.2008.07.004](https://doi.org/10.1016/j.psychres.2008.07.004)
- Reid, R. C., & Carpenter, B. N. (2009). Demoralization, Hypomanic Activation, and Disconstraint scores on MMPI-2 as significant predictors of hypersexual behavior. *Sexual Addiction and Compulsivity*, 16, 173–189. doi: [10.1080/10720160903202448](https://doi.org/10.1080/10720160903202448)
- Reid, R. C., & Carpenter, B. N. (2009). Exploring relationships of psychopathology in hypersexual patients using the MMPI-2. *Journal of Sex & Marital Therapy*, 35(4), 294–310. doi: [10.1080/00926230902851298](https://doi.org/10.1080/00926230902851298)
- Reid, R. C., Carpenter, B. N., & Draper, E. D. (2011). Disputing the notion of psychopathology among women married to hypersexual men using the MMPI-2-RF. *Journal of Sex & Marital Therapy*, 37, 45–55. doi: [10.1080/0092623X.2011.533585](https://doi.org/10.1080/0092623X.2011.533585)
- Sellbom, M., Bagby, R. M., Kushner, S., Quilty, L. C., & Ayearst, L. E. (2012). Diagnostic construct validity of the MMPI-2 Restructured Form (MMPI-2-RF) scale scores. *Assessment*, 19, 176–186. doi: [10.1177/1073191111428763](https://doi.org/10.1177/1073191111428763)
- Sellbom, M., Ben-Porath, Y. S., & Bagby, R. M. (2008). Personality and psychopathology: Mapping the MMPI-2 Restructured Clinical (RC) Scales onto the Five Factor Model of Personality. *Journal of Personality Disorders*, 22, 291–312. doi: [10.1521/pedi.2008.22.3.291](https://doi.org/10.1521/pedi.2008.22.3.291)
- Sellbom, M., Ben-Porath, Y. S., & Graham, J. R. (2006). Correlates of the MMPI-2 Restructured Clinical (RC) Scales in a college counseling setting. *Journal of Personality Assessment*, 86, 89–99. doi: [10.1207/s15327752jpa8601_10](https://doi.org/10.1207/s15327752jpa8601_10)
- Sellbom, M., Ben-Porath, Y. S., McNulty, J. L., Arbisi, P. A., & Graham, J. R. (2006). Elevation differences between MMPI-2 Clinical and Restructured Clinical (RC) Scales: Frequency, origins, and interpretative implications. *Assessment*, 13, 430–441. doi: [10.1177/1073191106293349](https://doi.org/10.1177/1073191106293349)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Sellbom, M., Graham, J. R., & Schenk, P. (2006). Incremental validity of the MMPI-2 Restructured Clinical (RC) Scales in a private practice sample. *Journal of Personality Assessment*, 86, 196–205. doi: [10.1207/s15327752jpa8602_09](https://doi.org/10.1207/s15327752jpa8602_09)
- Simms, L. J., Casillas, A., Clark, L. A., Watson, D., & Doebbeling, B. I. (2005). Psychometric evaluation of the Restructured Clinical Scales of the MMPI-2. *Psychological Assessment*, 17, 345–358. doi: [10.1037/1040-3590.17.3.345](https://doi.org/10.1037/1040-3590.17.3.345)
- Vachon, D. D., Sellbom, M., Ryder, A. G., Miller, J. D., & Bagby, R. M. (2009). A Five-Factor Model description of depressive personality disorder. *Journal of Personality Disorders*, 23, 447–465. doi: [10.1521/pedi.2009.23.5.447](https://doi.org/10.1521/pedi.2009.23.5.447)
- Van der Heijden, P. T., Egger, J. I. M., Rossi, G., & Derksen, J. J. L. (2012). Integrating psychopathology and personality disorders conceptualized by the MMPI-2-RF and the MCMI-III: A structural validity study. *Journal of Personality Assessment*, 94, 345–357. doi: [10.1080/00223891.2012.656861](https://doi.org/10.1080/00223891.2012.656861)
- Van der Heijden, P. T., Egger, J. I. M., Rossi, G., & Derksen, J. J. L. (in press). The MMPI-2-Restructured Form and the standard MMPI-2 Clinical scales in relation to DSM–IV. *European Journal of Psychological Assessment*.
- Wallace, A., & Liljequist, L. (2005). A comparison of the correlational structures and elevation patterns of the MMPI-2 Restructured Clinical (RC) and Clinical Scales. *Assessment*, 12, 290–294. doi: [10.1177/1073191105276250](https://doi.org/10.1177/1073191105276250)
- Watson, C., Quilty, L. C., Bagby, R. M. (2011). Differentiating Bipolar Disorder from Major Depressive Disorder using the MMPI-2-RF: A receiver operating characteristics (ROC) Analysis. *Journal of Psychopathology and Behavioral Assessment*, 33, 368–374. doi: [10.1007/s10862-010-9212-7](https://doi.org/10.1007/s10862-010-9212-7)
- Wolf, E. J., Miller, M. W., Orazem, R. J., Weierich, M. R., Castillo, D. T., Milford, J., Kaloupek, D. G., & Keane, T. M. (2008). The MMPI-2 Restructured Clinical Scales in the assessment of posttraumatic stress disorder and comorbid disorders. *Psychological Assessment*, 20, 327–340. doi: [10.1037/a0012948](https://doi.org/10.1037/a0012948)
- Wygant, D. B., & Fleming, K. P. (2008). Clinical utility of the MMPI-2 Restructured Clinical (RC) Scales in therapeutic assessment: A case study. *Journal of Personality Assessment*, 90, 110–118. doi: [10.1080/00223890701845112](https://doi.org/10.1080/00223890701845112)

Substance Abuse Treatment Settings

- Forbey, J. D., & Ben-Porath, Y. S. (2007). A comparison of the MMPI-2 Restructured Clinical (RC) and Clinical Scales in a substance abuse treatment sample. *Psychological Services*, 4, 46–58. doi: [10.1037/1541-1559.4.1.46](https://doi.org/10.1037/1541-1559.4.1.46)
- Monnot, M. J., Quirk, S. W., Hoerger, M., & Brewer, L. (2009). Racial bias in personality assessment: Using the MMPI-2 to predict psychiatric diagnoses of African American and Caucasian Chemical Dependency patients. *Psychological Assessment*, 21, 137–151. doi: [10.1037/a0015316](https://doi.org/10.1037/a0015316)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

Non-Clinical Settings

- Avdeyeva, T. V., Tellegen, A., & Ben-Porath, Y. S. (2011). Empirical correlates of low scores on MMPI-2/MMPI-2-RF Restructured Clinical Scales in a sample of university students. *Assessment, 19*, 388–393. doi: [10.1177/1073191111411675](https://doi.org/10.1177/1073191111411675)
- Baker, C. N., & Hoerger, M. (2012). Parental child-rearing strategies influence self-regulation, socio-emotional adjustment, and psychopathology in early adulthood: Evidence from a retrospective cohort study. *Personality and Individual Differences, 52*, 800–805. doi: [10.1016/j.paid.2011.12.034](https://doi.org/10.1016/j.paid.2011.12.034).
- Benitez, A., & Gunstad, J. (2012). Poor sleep quality diminishes cognitive functioning independent of depression and anxiety in healthy young adults. *The Clinical Neuropsychologist, 26*, 214–223. doi: [10.1080/13854046.2012.658439](https://doi.org/10.1080/13854046.2012.658439)
- Forbey, J. D., & Ben-Porath, Y. S. (2008). Empirical correlates of the MMPI-2 Restructured Clinical (RC) Scales in a non-clinical setting. *Journal of Personality Assessment, 90*, 136–141. doi: [10.1080/00223890701845161](https://doi.org/10.1080/00223890701845161)
- Forbey, J. D., & Lee, T. T. C. (2011). An exploration of the impact of invalid MMPI-2 protocols on collateral self-report measure scores. *Journal of Personality Assessment, 93*, 556–565. doi: [10.1080/00223891.2011.608757](https://doi.org/10.1080/00223891.2011.608757)
- Forbey, J. D., Lee, T. T. C., & Handel, R. W. (2010). Correlates of the MMPI-2-RF in a college setting. *Psychological Assessment, 22*, 737–744. doi: [10.1037/a0020645](https://doi.org/10.1037/a0020645)
- Hoerger, M., Quirk, S. W., & Weed, N. C. (2011). Development and validation of the Delaying Gratification Inventory. *Psychological Assessment, 23*, 725–738. doi: [10.1037/a0023286](https://doi.org/10.1037/a0023286)
- Ingram, P. B., Kelso, K. M., & McCord, D. M. (2011). Empirical correlates and expanded interpretation of the MMPI-2-RF Restructured Clinical Scale 3 (Cynicism). *Assessment, 18*, 95–101. doi: [10.1177/1073191110388147](https://doi.org/10.1177/1073191110388147)
- Kastner, R. M., & Sellbom, M. (2012). Hypersexuality in college students: The role of psychopathy. *Personality and Individual Differences, 53*, 644–649. doi: [10.1016/j.paid.2012.05.005](https://doi.org/10.1016/j.paid.2012.05.005)
- Lee, T. T. C., & Forbey, J. D. (2010). MMPI-2 correlates of sexual preoccupation as measured by the Sexuality Scale in a college setting. *Sexual Addiction and Compulsivity, 17*, 219–235. doi: [10.1080/10720162.2010.500500](https://doi.org/10.1080/10720162.2010.500500)
- Neal, T. M. S., & Sellbom, M. (2012). Examining the factor structure of the Hare Self-Report Psychopathy Scale. *Journal of Personality Assessment, 93*, 244–253. doi: [10.1080/00223891.2011.648294](https://doi.org/10.1080/00223891.2011.648294)
- Osberg, T. M., Haseley, E. N., & Kamas, M. M. (2008). The MMPI-2 Clinical Scales and Restructured Clinical (RC) Scales: Comparative psychometric properties and relative diagnostic efficiency in young adults. *Journal of Personality Assessment, 90*, 81–92. doi: [10.1080/00223890701693801](https://doi.org/10.1080/00223890701693801)

(continued)

MMPI-2-RF® Bibliography

Revised September 2012

- Sellbom, M., & Ben-Porath, Y. S. (2005). Mapping the MMPI-2 Restructured Clinical (RC) Scales onto normal personality traits: Evidence of construct validity. *Journal of Personality Assessment*, 85, 179–187. doi: [10.1207/s15327752jpa8502_10](https://doi.org/10.1207/s15327752jpa8502_10)
- Sellbom, M., Ben-Porath, Y. S., Lilienfeld, S. O., Patrick, C. J., & Graham, J. R. (2005). Assessing psychopathic personality traits with the MMPI-2. *Journal of Personality Assessment*, 85, 334–343. doi: [10.1207/s15327752jpa8503_10](https://doi.org/10.1207/s15327752jpa8503_10)
- Sellbom, M., Fischler, G. L., & Ben-Porath, Y. S. (2007). Identifying MMPI-2 predictors of police officer integrity and misconduct. *Criminal Justice and Behavior*, 34, 985–1004. doi: [10.1177/0093854807301224](https://doi.org/10.1177/0093854807301224)